

Evidence of small ferrimagnetic concentrations in mice (*Mus musculus*) livers and kidneys exposed to the urban dust : A reconnaissance study

Francisco Bautista, María E. Gonsebatt, Rubén Cejudo, Avto Goguitchaichvili, Ma. Carmen Delgado and Juan J. Morales

Received: August 15, 2017; accepted: November 09, 2017; published on line: January 01, 2018

Resumen

Investigaciones previas llevadas al cabo sobre el polvo y los suelos urbanos de la Ciudad de México revelaron una concentración relativamente alta de metales pesados debido a la contaminación atmosférica. Los análisis magnéticos detallados demostraron que las muestras de polvo urbano contienen una fase de magnetita neoformada procedente de la combustión de los vehículos. En este trabajo informamos los resultados de las medidas magnéticas realizadas en hígados y riñones de *Mus musculus* para evaluar si las partículas magnéticas pueden penetrar y acumularse en estos órganos vitales. Las medidas sistemáticas de la susceptibilidad magnética y las curvas de adquisición de magnetización remanente isotérmica atestiguan la existencia de pequeñas concentraciones de granos ferrimagnéticos de coercitividad baja / media (muy probablemente magnetita). Este estudio confirma que los minerales magnéticos finos pueden penetrar y llegar a hígados y riñones de ratones, en concentraciones relativamente pequeñas.

Palabras clave: Contaminación atmosférica, Ciudad de México, ratones, *Mus musculus*.

Abstract

Previous investigations carried out on Mexico City urban dust and soils revealed relatively high concentration of heavy metals due to the atmospheric pollution. Detailed magnetic analyzes demonstrated that the dust samples contain neo-formed magnetite phase coming from the vehicles combustion. Here, we report the results of magnetic measurements carried out on *Mus musculus* livers and kidneys in order to evaluate whether the magnetic particles may penetrate and accumulate in these organs. The systematic measurements of the magnetic susceptibility and isothermal remanent magnetization acquisition curves attest the existence of small concentrations of low/medium coercivity ferrimagnetic grains (most probably magnetite). This study confirms that fine magnetic minerals can penetrate and reach livers and kidneys of mice in relatively small concentrations.

Key words: Atmospheric pollution, Mexico City, mice, *Mus musculus*.

F. Bautista
Laboratorio Universitario de Geofísica Ambiental
Centro de Investigaciones en Geografía Ambiental
Universidad Nacional Autónoma de México
Morelia, Michoacán México
*Corresponding author: leptosol@ciga.unam.mx

M.E. Gonsebatt
Laboratorio de Toxicología Ambiental
Instituto de Biomédicas
Universidad Nacional Autónoma de México
Ciudad Universitaria, México.

R. Cejudo
A. Goguitchaichvili
C. Delgado
J. Morales
Laboratorio Universitario de Geofísica Ambiental
Instituto de Geofísica
Universidad Nacional Autónoma de México
Unidad Michoacán, México.

Introduction

Ten micrometer diameter (PM_{10}), or less, fine particles of anthropic origin contain magnetic minerals and traces of heavy metals that may be evidenced by magnetic methods as already reported in previous works (Qiong *et al.*, 2013; Ihl *et al.*, 2015; Cortés *et al.*, 2015, 2017). These studies have used different environmental samples (plant, soil, urban dust among others) to determine the relationship between the concentration of heavy metals (Cr, Cu, Pb, Zn, etc.) and some magnetic parameters (Wang, 2013; Cejudo *et al.*, 2015). The magnetic susceptibility (κ_f) and isothermal remanent magnetization values (IRM) up to 0.7 T have shown relatively good correlation with the concentration of the majority of heavy metals (Cd, Cr, Cu, Ni, Pb, V and Zn) (Evans and Heller, 2003; Qiao *et al.*, 2011; Bautista *et al.*, 2014; Cejudo *et al.*, 2015). The poor air quality in mega cities is mainly related with gas emission by cars, as they systematically emit fine particles containing both magnetic materials and heavy metals. These fine particles accumulate on the surface of the soil, floors (concrete, pavement, paving stones, etc.) and constitute the urban dust (Bautista *et al.*, 2011; Cejudo *et al.*, 2015; Sánchez-Duque *et al.*, 2015; INECC, 2011; 2016).

The population is naturally exposed to urban dust that may penetrate into the human body through oral ingestion, breathing and skin contact (Baroli *et al.*, 2007; Kwon *et al.*, 2008). In humans, the particles with a size of 2.5 microns in diameter can enter through the respiratory tract to the region of the pulmonary alveoli and ultra-fine particles (less than 1 micron in diameter) can even penetrate into the circulatory system (Forsman and Hogstedt, 1995).

Digestive system has an important role in nutrient acquisition within the stomach and intestines where food and liquids are mixed with digestive juices to transform it into smaller molecules, which are absorbed by the intestinal walls and driven by the blood and the hepatic vein to the liver (Townsend *et al.*, 2014). After necessary nutrients have been taken by the body, the residual waste is sent to the blood through the kidneys, where the filtration is achieved to remove most toxins and other substances (pesticides, drugs and food additives). The wastes removal occurs in tiny units inside the kidneys called nephrons, which act as filters where waste and excess water are combined and become urine (Guyton and Hall, 2011).

The main objective of this study is to reveal (if any) and quantify the presence of magnetic minerals in livers and kidneys of mice exposed to urban dust.

Magnetic experiments

In order to know whether the urban dust potentially penetrates in mice vital organs, a specific experimental procedure was carried out using three types of urban dust from Mexico City with different (contrasting) values of mass specific susceptibility and isothermal remanent magnetization at 0.7 T. (DIL, DIIL, DIIL) and a control (CL).

The urban dust was dried in the shade and sieved to 2 mm to remove the coarse-grained fraction from the gravel. The material was put into 8 cc plastic boxes, ideal for magnetic measurements. Magnetic susceptibility (κ_f) was measured at low frequency (0.46 kHz) with a MS2B Bartington and isothermal remanent magnetization (IRM) up to 0.7 T obtained at room temperature using a pulse magnetizer IM-10 (ASC Scientific).

The values of mass specific susceptibility ($\chi_{lf} = \kappa_f/\rho$, density) and $IRM_{0.7T}$ (at 0.7 T) were recorded for each urban dust samples. The urban dust was classified according to the measurements of mass specific susceptibility and isothermal remanent magnetization in low, medium and high values. A mixture of 1000 mL saline solution and 1 g of each urban dust was prepared. The mice were dosed with 16 ml per day during a total of twelve days.

Fifteen mice (*Mus musculus*) were divided into three lots; each lot was assigned a saline solution prepared with urban dust. The first lot of mice (Group I) was dosed with a urban dust solution having low values of $IRM_{0.7T}$; the second lot (Group II) was dosed with a urban dust solution having medium values of $IRM_{0.7T}$ and the third lot of mice (Group III) was dosed with a urban dust solution with high values of $IRM_{0.7T}$. A control lot of five mice was also used to which a saline solution without urban dust was applied.

The time and dosed amount of the solution was the same for each lot during twelve days. Twenty samples of liver and kidney were obtained after mice sacrifice. The organs were separated immediately after sacrifice, and frozen at -70°C . The organ samples were preserved at 4°C in 8 cc plastic boxes for magnetic measurements.

Once the saturation was achieved at 0.7 T, a magnetic field of 0.2 T was applied in the opposite direction to measure the $IRM_{-0.2}$ value and to determine the ratio S_{-200} ($S_{-200} = IRM_{-0.2} / IRM_{0.7}$). S_{-200} reflects the relative proportion of high coercivity minerals (values less than 0.75) and low coercivity minerals (values between 0.75 and 1.0) (Thompson and Oldfield, 1986; Bloemendal *et al.*, 1992; Gubbins and Herrero-Bervera, 2007). All magnetizations were measured with a Molspin (ASC Scientific) and JR6 (AGICO) spinner magnetometers.

A simple variance analysis was performed between the groups of mice dosed with saline solution and the magnetic measurements (k_{if} , NRM, $IRM_{0.7}$, $IRM_{0.2}$ and S_{-200}). A factorial variance analysis was carried out using as factor to organs (liver and kidney) and groups of urban dust. A medium comparison analysis was carried out (Statgraphic, 2007).

Main results, discussion and concluding remarks

The three urban dust samples exhibited different values of mass specific magnetic (χ_{if}) and isothermal remanent magnetization ($IRM_{0.7T}$); urban dust I (group DIL) is characterized with relatively low value of $IRM_{0.7T}$ (27.7 $mAm^2 kg^{-1}$); urban dust II (Group DIIL) with a medium value of $IRM_{0.7T}$ (49.0 $mAm^2 kg^{-1}$) while the urban dust III (group DIIL) yielded the highest value of $IRM_{0.7T}$ (95.2 $mAm^2 kg^{-1}$) (Table I).

Table 1. Results of magnetic measurements of three urban dust groups.

Sample	χ_{if} $\mu m^3 kg^{-1}$	$IRM_{0.7T}$ $mAm^2 kg^{-1}$
Dust I	2.1	27.7
Dust II	2.8	49.0
Dust III	3.3	95.2

After twelve days dosing, 20 samples of liver and kidney were respectively obtained. The average mass for all liver sample was 0.5 g and kidney reached 0.3 g. The magnetic parameters were not normalized by density, since it was impossible to determine the sample density without affecting the integrity of the material.

The negative initial susceptibility values obtained in liver samples are very likely due to the high concentration of diamagnetic material (mainly water and organic matter) (Table II). The natural remanent magnetization (NRM) yielded values between 0.0 to 0.1 $mA m^{-1}$; the

$IRM_{0.7}$ provided values between 0.6 to 2.1 $mA m^{-1}$; the $IRM_{0.2}$ provided values between 0.6 to 2.1 $mA m^{-1}$ and the S_{-200} ratio showed values higher than 0.8 (Figure 1 and Table II).

The kidney samples showed magnetic susceptibility values between -9.1 to 5.0×10^{-6} (SI); NRM values between 0.0 to 0.1 $mA m^{-1}$; $IRM_{0.7}$ values between 0.6 to 2.0 $mA m^{-1}$; the $IRM_{0.2}$ showed values between 0.6 and 2.0 $mA m^{-1}$ and the S_{-200} ratio showed the higher values between 0 and 2.3. Values of S_{-200} close to 1.0 indicate high proportion of low coercivity minerals. In a single case however, R4S1 kidney sample showed a value less than 0.7 indicative of high coercivity minerals (Table II, Figure 2).




Figure 1. Box plot of magnetic susceptibility values and isothermal remanent magnetization at 0.7 T.

The variance statistical analysis yielded significant differences between the values obtained from liver samples. The highest values correspond to the DIIL treatment with higher concentration of ferromagnetic minerals (Figure 1). In case of $IRM_{0.7}$ parameters, although significant changes were also detected, the values from control samples resulted relatively high most probably due to simple contamination.

The kidney samples associated with the high magnetic concentration urban dusts showed higher values of both susceptibility and $IRM_{0.7}$ as expected (Figure 2).

Table 2. Magnetic susceptibility and isothermal remanent magnetization values at 0.7 T for liver and kidney samples.

Sample	Weight g	K_{if} $\times 10^{-6}(\text{SI})$	NRM mA m^{-1}	$\text{IRM}_{0.7\text{T}}$ mA m^{-1}	$\text{IRM}_{-0.2\text{T}}$ mA m^{-1}	S_{-200}	S_{-25}
Liver samples							
Min	0.4	-10.5	0.0	0.6	0.6	0.8	0.85
Max	0.6	6.0	0.1	2.1	2.1	1.0	1.00
Mean	0.5	0.0	0.1	1.2	1.2	0.9	0.93
Kidney samples							
Min	0.2	-9.1	0.0	0.6	0.6	0.1	0.00
Max	0.4	5.0	0.1	2.0	2.0	2.3	0.70
Mean	0.3	-1.8	0.0	1.1	1.1	1.0	0.2

Magnetic parameters such as κ_{if} and $\text{IRM}_{0.7}$ showed that they could be used for the identification of magnetic signal in mouse organs by ingestion of urban dust. The natural remanent magnetization measurements were not adequate to identify the intake of magnetic particles in mice at low exposure.

The liver samples showed the highest concentrations of magnetic particles compared to the kidney samples. It follows that, at low exposure times to urban dust, damage could be detected first in the liver.

The IRM acquisition curves in liver samples exhibited the following behavior: between 0 to 100 mT fields, the remanent magnetization increases proportionally with the applied field. For magnetic fields between 100 to 200 mT, the growth of the magnetization stops and the saturation is reached. Liver samples of Group I showed relatively low values of saturation remanent magnetization ($\text{IRM}_{0.7} = 0.75 \text{ mA m}^{-1}$), while higher $\text{IRM}_{0.7}$ values were reported for group III ($\text{IRM}_{0.7} = 2.2 \text{ mA m}^{-1}$) (Figure 2 and 3).

High IRM values of 1.5 mA m^{-1} were recorded for kidney samples of group III, whereas samples of group II and I exhibited almost the same value of $\text{IRM}_{0.7}$ (1.2 mA m^{-1} , Figure 3).

The liver and kidney samples showed values of S_{-200} close to 1.0 indicating high proportions of low coercivity minerals (LCM) and low proportion of high coercivity mineral (HCM) (Figure 4).

The results obtained in this study showed that the liver and kidney of mice of *Mus musculus* species present significant magnetic response to the applied fields. The group of mice dosed with the highest concentrations of magnetic particles (dust III) yielded a higher concentration of magnetic minerals in their liver and kidney and vice versa. These experiments also indicated that low coercivity ferrimagnetic minerals (possibly magnetite) are present in the liver and kidney of mice.

On the other hand, magnetic particles are being used in biomedical research because




Figure 2. Magnetic susceptibility and isothermal remanent magnetization values at 0.7 T for kidney samples.

they can be carriers of drugs to attack malignant tumors (Chao *et al.*, 2012) and to trap elements such as heavy metals and toxic substances in the blood (Ramanujan, 2009), and other apps (Li *et al.*, 2016)..It is also known that magnetic particles in high concentration can cause cytotoxicity, DNA damage, oxidative stress, epigenetic events and inflammatory processes (Sing *et al.*, 2010); Furthermore, it can accumulate in organs such as liver, brain,

spleen and lungs subsequent to inhalation and penetration through hair follicles (Baroli *et al.*, 2007; Kwon *et al.*, 2008). However, little research has been done on the use of magnetic particles as indicators of contamination in animal tissues.

The distribution, transformations and biotransformations of the magnetic particles in the body are scarcely known (Levy *et al.*, 2011).




Figure 3. Isothermal remanent magnetization acquisition curves for liver and kidney *Mus musculus* samples.




Figure 4. Relationship between S_{-200} and measures of $IRM_{0.7}$ for all liver (circles) and kidney (square) samples.

In addition, when the research is done with environmental samples, such as urban dust (mixture of particles in both size and chemical composition) very few studies are found.

These type of studies are relevant also for human health, showing that adults ingest approximately 50 mg per day of ambient dust and soil, while infants or children ingest twice as much (100-110 mg per day) (US EPA, 2011).

The main finding of the present study is that the liver was the organ with the highest magnetic signal. The kidney also achieved a magnetic signal but of smaller magnitude as compared with the liver.

This reconnaissance study confirms that fine magnetic minerals may be absorbed and stored by mice livers and kidneys in relatively small concentrations but also in small exposure time. The same mechanism may be speculated for the human vital organs. However, more experiments are required considering: a) a higher mice population; b) longer exposure to urban dust; c), the introduction of control lots; and e) the identification of the magnetic particles.

Acknowledgments

To Secretaria de Ciencias, Tecnología e Innovación de la Ciudad de México for the financial support of the project SECITI/051/2016. We thank the National Science and Technology Council (CONACYT) for the financial support CB-2011-01-169915. Authors would like to thank Hilda Rivas for technical support.

References

- Baroli B, Ennas MG, Loffredo F, Isola M, Pinna R, Lopez-Quintela MA. 2007. Penetration of metallic nanoparticles in human full-thickness skin. *J Invest Dermatol.* 127:1701-12.
- Bautista F, J.L. Palacio H. Delfín y (Eds.). 2011. Técnicas de muestreo para el estudio del manejo de recursos naturales y el cuidado del ambiente. Centro de Investigaciones en Geografía Ambiental. Universidad Nacional Autónoma de México. México D.F. 790 pp.
- Bautista F, Cejudo R., Aguilar B., Goguichaisvili A., 2014. El potencial del magnetismo en la clasificación de suelos: una revisión. *Bull. Soc. Geol. Mex.* 66, 2, 365-376.

- Bloemendal, J., King, J.W., Hall, F.R., and Doh, S.-J., 1992. Rock magnetism of Late Neogene and Pleistocene deep-sea sediments: relationship to sediment source, diagenetic processes, and sediment lithology. *Journal of Geophysical Research*, 97: 4361–4375
- Cejudo R., Bautista F., Quintana P., Delgado A., Aguilar A., Gogitchaishvili A., Morales J., 2015. Correlación entre elementos potencialmente tóxicos y propiedades magnéticas en suelos de la Ciudad de México para la identificación de sitios contaminados: definición de umbrales magnéticos. *Rev. Mex. Cienc. Geol.* 32, 1, 50-61.
- Cortés J.L.; F. Bautista; P. Quintana; D. Aguilar y A. Gogichaishvili. 2015. El color del polvo urbano como indicador de contaminación por elementos potencialmente tóxicos: El caso de Ensenada, Baja California. *Revista Chapingo Serie Ciencias Forestales y del Ambiente.* 20 (2): 255-266. Doi: dx.doi.org/10.5154/r.rchscfa.2015.02.003.
- Cortés J.L.; F. Bautista; P. Quintana; y A. Gogichaishvili. 2017. Distribución espacial de los metales pesados en polvos urbanos de la ciudad de Ensenada, Baja California, México. *Revista Chapingo Serie Ciencias Forestales y del Ambiente.* 23(1): 235-248.
- Chao, X, Zhang Z, Guo L, Zhu J, Peng M., Vermorken A., Van de Ven W, Chen C. and Cui Y. 2012. A novel magnetic nanoparticle drug carrier for enhanced cancer chemotherapy. *PLoS one* 7(10), e40388, <https://doi.org/10.1371/journal.pone.0040388>
- Evans M.E., Heller F., 2003, *Environmental Magnetism- Principles and Applications of Enviromagnetics.* Int. Geophysics Ser., 86, Academic press, Amsterdam, 293 pp.
- Forsman M and Hogstedt P 1995. Total lung dust inverse solutions in single plane SQUID magnetopneumography” *Biomagnetism: Fundamental Research and Clinical Applications* (Amsterdam: Elsevier) pp 793–7.
- Guyton A., Hall J., 2011. *Tratado de fisiología médica.* Barcelona España, Elsevier. 12 edición, 1083 pp.
- Gubbins D. and HerreroBervera E., 2007. *Encyclopedia of Geomagnetism and Paleomagnetism,* Springer, Heidelberg, 1072 p.
- Kwon JT, Hwang SK, Jin H, Kim DS, Minai-Tehrani A, Yoon HJ, Han DY, Kang YW, Yoon BI, Lee JK, Cho MH. 2008. Body distribution of inhaled fluorescent magnetic nanoparticles in the mice. *J Occup Health;* 50:1–6.
- Ihl T., F. Bautista, R. Cejudo*, C., Delgado*, AvtoGoguichaishvili, P Quitana y D. Aguilar. 2015. Concentration of toxic elements in topsoils of Metropolitan area of Mexico City: A spatial analysis using Ordinary Kriging and Indicator Kriging. *Revista Internacional de Contaminación Ambiental.* 31: 42-62.
- INECC. Instituto Nacional de Ecología, 2011, *Guía metodológica para la estimación de emisiones de PM2.5,* INE-Semarnat, México D.F. 105 pp.
- INECC. Instituto Nacional de Ecología, 2016, *Metales pesados,* <http://www.inecc.gob.mx/sqre-temas/763-aqre-metales>, consultado en enero 2016.
- Kwon JT, Hwang SK, Jin H, Kim DS, Minai-Tehrani A, Yoon HJ, Han DY, Kang YW, Yoon BI, Lee JK, Cho MH. 2008. Body distribution of inhaled fluorescent magnetic nanoparticles in the mice. *J. Occup. Health;*50:1–6.
- Levy M., Wilhelm C., Luciani N., Deveaux V., Gendron F., Luciani A., Devaud M. and Gazeau F., 2011. Nanomagnetism reveals the intracellular clustering of iron oxide nanoparticles. *Nanoscale,* 3, 4402.
- Li XM, Wei JR, Aifantis KE, Fan Y, Feng Q, Cui F, Watar F. 2016. Current investigations into magnetic nanoparticles for biomedical applications. *J. Biomed. Mater Res. A.* 104(5):1285–1296.
- Qiao Q., Zhang C., Huang B., Piper J., 2011. Evaluating the environmental quality impact of the 2008 Beijing Olympic Games: magnetic monitoring of street dust in Beijing Olympic Park, *Geophys. J. Int.,*187, 1222–1236.
- Qiong W., Xiao-Hui B., Jian-Hui W. Yu-Fen Z., Yin-Chang F. 2013. Heavy metals in urban ambient PM10 and soil background in eight cities around China, *Environmental Monitoring and Assessment,* 185:1473–1482.
- Ramanujan R.V. 2009. Chapter 17 Magnetic Particles for Biomedical Applications. En: R. Narayan (ed.), *Biomedical Materials.* Springer. 467-491 pp. DOI 10.1007/978-0-387-84872-3 17.

Sánchez-Duque, A.*, F. Bautista, A. Gogichaishvili, R. Cejudo-Ruiz, J. Reyes-López, F. Solís-Domínguez y J. Morales-Contreras. 2015. Evaluación de la contaminación ambiental a partir del aumento magnético en polvos urbanos - Caso de estudio para la ciudad de Mexicali, México. *Revista Mexicana de Ciencias Geológicas*. 32:501-513.

Singh, N., Jenkins, G. J. S., Asadi, R., and Doak, S. H. 2010. Potential toxicity of superparamagnetic iron oxide nanoparticles (SPION). *Nano Reviews*, 1, 10.3402/nano.v1i0.5358. <http://doi.org/10.3402/nano.v1i0.5358>

Statgraphic 2007, ANOVA Simple, 1-24, <http://www.statgraphics.net/wp-content/uploads/2011/12/tutoriales/ANOVA%20Simple.pdf> consulted on february 3, 2016.

Thompson, R., Oldfield, F., 1986, *Environmental Magnetism*: London, Allen and Unwin, 227 pp.

Townsend C., Beachamp R., Marck B., Mattox K., 2014, *Sabiston Cirugía General y del Aparato Digestivo*, Barcelona España, Elsevier, 576 p.

U.S. EPA. Exposure Factors Handbook 2011 Edition (Final). U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-09/052F, 2011.

Wang X. 2013. Assessment of heavy metal pollution in Xuzhou urban topsoils by magnetic susceptibility measurements. *Journal of Applied Geophysics*, 92, 76-83.