

HIGH PRECISION MEASUREMENTS OF ABSOLUTE GRAVITY IN MÉXICO: THE JALISCO BLOCK CHANGES IN GRAVITY TRIGGERED BY DISTANT EARTHQUAKES

Alfredo Esparza^{1,2}, Jorge Arzate^{*3}, Ludger Timmen⁴, Jason Silliker⁵ and Manuel Schilling⁴

Received: August 27, 2019; accepted: March 25; published online: July 1, 2020

RESUMEN

En este trabajo se reportan los resultados de 16 mediciones de gravedad absoluta (GA) utilizando dos gravímetros de caída libre, el FG5X-220 de la Universidad Leibniz de Hannover (LUH) y el FG5X-252 del Centro Nacional de Metrología (CENAM). Previo al establecimiento de nuevas estaciones gravimétricas de primer orden y a las campañas de adquisición en el Bloque de Jalisco (BJ), se llevó a cabo la certificación del gravímetro FG5X-252 a partir de dos comparaciones instrumentales; la primera con el FG5X-220 de LUH, el cual es un instrumento certificado de larga estabilidad y repetibilidad debajo de los 2 μGal , y la segunda a través de una comparación internacional con otros 13 instrumentos en las instalaciones de la NOAA en Table Mountain, Colorado, USA.

Las campañas de medición en el BJ se realizaron durante la estación seca (Feb/Mar) los años 2016 y 2018, e incluyó las estaciones de Chamela (CHA), Guadalajara (UGG), Manzanillo (MAN), Puerto Vallarta (UGP) y Tepic (TEP), ésta última establecida como nueva referencia en el norte del bloque. Los resultados obtenidos de estas dos campañas en el BJ fueron comparados con los valores medidos en 1996 por la NOAA en esta misma región del oeste de México. Los desplazamientos verticales observados en el lapso de dos años en las estaciones CHA (+22.7 cm), UGG (+44.3 cm) y MAN (+54.6 cm) supera sustancialmente el promedio anual (2.8 cm, 4.2 cm y 3.6 cm respectivamente) con respecto a las mediciones de GA de 1996. En el mismo periodo, la estación UGP subsidió 8.5 cm mientras que TEP permaneció muy estable (-0.25 cm).

En septiembre de 2017 ocurrieron dos grandes sismos de magnitudes 8.2 y 7.1, con epicentros en la costa de Chiapas (07/sep/2017) y en el Estado de Puebla (19/sep/2017), que fueron registrados en algunas de las estaciones GPS de la red UNAVCO en los sitios MAN, UGG y CHA a pesar de que la fuente sísmica más cercana se localiza a más de 500 km de distancia. El análisis de los datos adquiridos y otros datos geofísicos disponibles apoyan la hipótesis de que el sismo con epicentro en Puebla disparó la subducción asísmica de una porción de la placa de Rivera (PR), lo que a su vez produjo el levantamiento en dichas estaciones. Concluimos que la subducción asísmica en la zona ocurre debido a la subducción de una corteza oceánica hidratada que arrastra un importante espesor de sedimentos marinos, lo que genera una interfaz lubricada.

*Corresponding author
arzatej@geociencias.unam.mx

¹Centro Nacional de Metrología (CENAM) km 4.5
Carretera a Los Cués, El Marqués, Qro., CP 76246, México

²Posgrado en Ciencias de la Tierra, Campus UNAM-
Juriquilla, Blvd. Juriquilla #3001, CP 76230, Querétaro,
México

³Centro de Geociencias, Campus UNAM-Juriquilla,
Blvd. Juriquilla #3001, CP 76230, Querétaro,
México

⁴Leibniz Universität Hannover (LUH), Schneiderberg
50, 30167 Hannover, Germany

⁵Natural Resources Canada (NRC/RNC),
Ottawa, Canada

Palabras clave: g-Absoluta, comparación internacional, desplazamientos verticales, Bloque de Jalisco, subducción asísmica

ABSTRACT

We report the results of 16 Absolute Gravity (AG) measurements distributed central and western Mexico employing two free-fall gravity instruments; the FG5X-220 of Leibniz Universität Hannover (LUH), and the FG5X-252 instrument of the Centro Nacional de Metrología (CENAM). Previous to the setup of new stations and acquisition campaigns, the FG5X-252 was certified in two steps, first a mutual comparison with the reference gravimeter FG5X-220, which have a long range stability below 2 μGal , and later through an international comparison at NOAA's Table Mountain, Col. facilities with 13 other instruments of different countries.

The acquisition campaigns in the Jalisco Block (JB) took place during the dry season of 2016 and 2018, which included AG stations in Chamela (CHA), Guadalajara (AGG), Manzanillo (MAN), Puerto Vallarta (UGP), and Tepic (TEP); the later established new reference station in the north of the JB in 2016. The results obtained from the 2016 and 2018 field campaigns in the JB were compared with 1996 AG data acquired by NOAA at the same sites established. The observed vertical displacements in the two years period at stations CHA (+22.7 cm), UGG (+44.3 cm) and MAN (+54.6 cm) overcomes substantially the annual average (2.8 cm, 4.2 cm y 3.6 cm respectively) from the difference of the AG measurements 2016-1996. In the same period the UGP station subsided 8.5 cm, while station TEP remained quite stable (-0.25).

In September 2017 two large earthquakes of magnitudes 8.2 and 7.1 occurred in the coast of Chiapas (07/Nov/2017) and in the State of Puebla (19/Nov/2017), which were recorded at some of the UNAVCO's GPS stations, namely MAN, UGG and CHA even though the nearest seismic source was located more than 500 km to the east. The analysis of our results in combination with other geophysical data support the hypothesis that the earthquake with epicenter in Puebla triggered the aseismic subduction of a segment of the Ribera Plate (RP), which in turn uplifted the stations above mentioned. We conclude that the aseismic subduction in this region is facilitated by a wet oceanic crust that carries important amounts of marine sediments, producing a lubricated interface between oceanic RP and the overriding JP.

Key words: Absolute gravity, international comparison, vertical displacements, Jalisco Block, aseismic subduction

INTRODUCTION

The first precise gravity measurements in Mexico were carried out in the early 1970s (Bureau Gravimétrique International, BGI). Then, the g-values were derived from relative gravity surveys with ties to absolute determined points promoted by the Inter-American Geodetic Survey (IAGS). The accuracy of most of the 260 reference gravity stations (see Figure 1) is in the order of 0.1 mGal, however, the precise location of the measured gravity stations is not well known. In 1996 a first order absolute gravimetric survey in western Mexico was carried out aimed at

geodynamic investigations, led by the American National Oceanic and Atmospheric Administration (NOAA) with participation of the Instituto de Geofísica-UNAM. Their field campaign totaled the setting of 9 stations established by Daniel Winester of NOAA, using the NSF-funded absolute gravimeter FG5-111, and owned by the University of Colorado at Boulder (Roger Bilham, pers. com.). Their measurement included four stations located within the Jalisco Block, namely Chamela (CHA), Guadalajara (UGG), Manzanillo (MAN) and Puerto Vallarta (UGP) stations, other station at the basement of the Instituto de Geofísica-UNAM in Mexico City (IGU), and 4 more stations in the cities of Acapulco, Chilpancingo, Pinotepa and Taxco. However, the later four were not visited because the reference marks were destroyed.




Figure 1. Relative gravity stations measured in the 60's and early 70's in Mexico as archived by the Bureau Gravimétrique Internationale or BGI (<http://bgi.omp.obs-mip.fr/data-products/Gravity-Databases/Reference-Gravity-Stations>).

As part of this work, the Jalisco Block stations (UGP, CHA, MAN, UGG, shown in Figure 2) and the reference UNAM-CU station (IGU), were measured at the same benchmark points established during the NOAA's winter field campaign of 1996 in Mexico. Our measurements were also done during the winter of 2016 and repeated two years later in 2018 during the same time of the year (Feb-Mar). This was intentionally planned to prevent unwanted variations of g due to water table rise. In addition, we measured a new station in Tepic (TEP), north of the JB, and updated the INEGI main reference site (AGS) in Aguascalientes city. The remaining two stations (CGE and CNM) were established in August 2015 with the FG5X-220 absolute gravity meter and were measured again in February 2016 with the FG5X-220 and in March 2018 with the FG5X-252. The CGE station is located at Centro de Geociencias (CGEO-UNAM), and station CNM at the Gravity Laboratory of Centro Nacional de Metrología (CENAM), both in Querétaro. The 2016 AG campaign was conducted by a combined team from the University of Leibniz, Hannover (LUH), CENAM, and UNAM, and the 2018 measuring campaign by a combined team of CENAM, UNAM and NRC of Canada.




Figure 2. Absolute gravity (AG) stations measured during the first (2016) and second (2018) campaigns (red and yellow circles). Four of the five stations in the Jalisco Block (red circles) were established in 1996 (NOAA) and were used as reference for comparison. UNAM first order gravity station in Mexico City was also established in 1996. We setup the control Tepic (TEP) station, north of the JB, and updated the INEGI main reference site (AGS) in Aguascalientes city. The remaining two stations (CGE and CNM) were established in August 2015 and were measured again in February 2016 and in March 2018. The CGE station is located at Centro de Geociencias (CGEO-UNAM), and station CNM at the Gravity Laboratory of Centro Nacional de Metrología (CENAM), both in Querétaro. Green arrows indicate convergence direction of RP, and shaded grey zones represent the extension of the trans Mexican volcanic belt (TMVB). Lighter grey area delineates younger vulcanism. The Middle America Trench (MAT) is shown with brown line. The Rivera and Tamayo transform fault zones (RFZ and TFZ respectively) are highlighted with a NW-SE.

COMPARISONS AND CERTIFICATION OF THE FG5X-252

To start using a new absolute gravimeter with certainty is always necessary to know about the stability and repeatability of the g measurements that yields. The international comparison of CENAM's FG5X-252 absolute gravity meter acquired in December 2015, was done in two steps. First, in February 2016, a comparison with the LUH's international reference FG5X-220 gravity meter (Jiang *et al.*, 2012; Francis *et al.*, 2015; Timmen *et al.*, 2015; Schilling and Timmen, 2016), which produced a difference of $7.98 \pm 2.5 \mu\text{Gal}$ after corrections (Figure 3a). The FG5X-220 is a well-defined instrument that has participated in multiple international comparisons, including the last CCM.G.K² key comparison (Francis *et al.*, 2015) at the Underground Laboratory for Geodynamics in Walferdange, Luxembourg (2013). During the latest Regional key comparison

of Absolute Gravimeters, EURAMET.M.G-K2 and Pilot Study at the University of Luxembourg in Belval in November 2015, an expanded uncertainty of 5 μGal was derived for the Hannover FG5X-220 instrument, and its long term stability and repeatability is below 2 μGal (Schilling and Timmen, 2016). A second comparison of the absolute gravity meter was done in October of 2017 at the international comparison in Table Mountain in Boulder, CO (https://www.bipm.org/utils/common/pdf/final_reports/M/G-K1/SIM.M.G-K1.pdf), where a group of 13 instruments from 10 different countries participated. Figure 3b show the results of that comparison, where it is observed that the FG5X-252 was among the more stable. The first instrumental comparison was done before the 2016 field campaign, while the international certification at Table Mountain was made before the 2018 field campaign.




Figure 3. a) Comparison between FG5X-220 and FG5X-252 at the micro-Gravimetry Laboratory of CENAM in February 2016; b) results of the international comparison of 12 absolute gravity (AG) meters at Table Rock (NOAA, Boulder, CO) in Oct., 2017; c) aspects of FG5X-252 AG meter comparison at CENAM (left) and at Table Rock laboratory (right). The transportation of an AG gravimeter type FG5X (Micro-g®) involves the packing and the carrying of eight boxes worth ~ 250 kg (center).

APPLIED CORRECTIONS

To deduce the g-value from the 3,000 time/distance raw data pairs (<http://www.microglacoste.com/>), the applied equation of motion is given by:

$$x_i = x_0 + v_0 \left(\tau_i + \frac{\gamma}{6} \tau_i^3 \right) + \frac{g_0}{2} \left(\tau_i^2 + \frac{\gamma}{12} \tau_i^4 \right) \quad (1),$$

where x_0 , v_0 and g_0 stand for the initial position, the initial velocity and the gravity value at $t=0$ respectively. Here γ is the vertical gravity gradient ($\delta g/\delta h$) at the measuring site, (t_i, x_i) are the i -th time and interference derived position of the free-fall test body during a drop, and τ_i is the corrected time for the finite speed of light, given by (Nagorny, et. al 2010, Rothleitner, et. al 2014)

$$\tau_i = t_i - \frac{(x_i - x_0)}{c} \quad (2).$$

The individual g -values are corrected for three global geophysical effects: the gravitational Earth tides, the varying gravity attraction and loading effect due to mass redistributions in the atmosphere during measurements, and the change in the centrifugal acceleration due to polar motion. Here, the tidal variations have been predicted using Timmen and Wenzel (1995) algorithm, which computes the gravity tide parameters from gravity tide amplitudes of the Tamura's (1987) tidal potential. The obtained parameters include the corrections due to the solid Earth tides and the loading and attraction effect from the ocean tides. To carry out the reductions due to atmospheric variations we used the equation (IAG, 1983).

$$C_p = A (P_0 - P_n), \quad (3),$$

that expresses the barometric pressure C_p (in μGal) in terms of the observed atmospheric pressure P_0 (in hPa), and the nominal pressure P_n at the site. In this equation A is the barometric admittance factor ($-0.3 \mu\text{Gal}/\text{mBar}$). The nominal pressure P_n at a site located at an elevation h_m is estimated by the empiric equation (US Standard Atmosphere, 1976. (<https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19770009539.pdf>):

$$P_n = 1013.25 * (1 - 0.0065 \frac{h_m}{288.15})^{5.2559} \quad (4).$$

The variation in gravity due to polar motion (δ_{gpm} in μGal) during the measuring periods can be accounted for applying the equation (Torge and Müller; 2012)

$$\delta_{gpm} = -1.164\omega^2 a \text{sen}^2(2\varphi)(x \cos \lambda - y \sin \lambda) \quad (5),$$

here ω is the Earth's angular rotational velocity (rad/s), a is the equatorial radius (semi-major axis) of the reference ellipsoid (m), φ is the latitude of the site, λ the longitude, and x, y the pole coordinates (rad) as provided by the International Earth Rotation and Reference Systems Service (IERS) (<http://www.iers.org/IERS/EN/EarthOrientationData>). Polar motion and air pressure corrections were calculated using the g -software from Micro- g Lacoste® (<http://www.microglacoste.com/fg5x.php>). In order to compare our results with those measured during the 1996 acquisition campaign with the FG5-111 gravity meter, all measured g

values were transferred to the floor mark using a nominal sensor height of 1.30 m. The local gravity gradients used here were the same measured by the NOAA team in 1996; at those stations where the gradient was not available, we used the reference value of $-3 \mu\text{Gal}/\text{cm}$ (TEP, AGS, CNM, CGE).

FIELD MEASUREMENTS

During the field campaign of March 2016, the gravity meter FG5X-252 stayed at the CENAM Gravity Laboratory while the FG5X-220 LUH instrument was used for the measurements elsewhere. A complete FG5X instrument weights about 320 kg and is transported in 6 large boxes (Figure 3c) adding up to a packing volume of 1.5 m^3 approximately. The space floor requirements for installation are about 3 m^2 and a stable electric power supply of 110-240 VAC at 50-60 Hz. Nominal power requirements are 500 W sustained for at least 18 hours. The setup of the instrument takes about 1 to 2 hours for full operation it requires time to stabilize to room temperature ($15\text{-}25^\circ\text{C}$). Although the FG5Xs have a worldwide dynamic range, they have a limited operating temperature range, which is necessary to keep as stable as possible during a measurement. During the second field campaign the temperature conditions in the coastal region climbed above 35°C during the day but settled during the night above the low twenties, particularly when measuring in Chamela (CHA) and CUNIVO (MAN). We observed that a temperature variation of $5\text{-}10^\circ\text{C}$ degree overnight can lead to total failure of the automatically running observations, or to a loss in accuracy of more than $2 \mu\text{Gal}/\text{deg}$. Once settled and stabilized, the gravimeter is programed to acquire data series over night until next late morning assuring at least one semi-diurnal tidal period for an adequate correction of data. A full station measurement typically consists of up to 3,000 single free-fall drops events which are corrected for Earth’s tides, atmospheric mass redistributions and polar motion. To transfer the g -result from the FG5X sensor height to the floor level, the vertical gravity gradient $\delta g/\delta h$ was taken as $3.000 \mu\text{Gal}/\text{cm}$ at those stations where the gradient was not available. Table 1 shows the summary of results from this and the previous acquisition campaigns.

Table 1. Location, coordinates, and corrected absolute gravity g values corresponding to the NOAA’s 1996 campaign, and results of the first (2016) and second (2018) field campaigns as part of this work.

STN	LOCATION	Lat	Long	Z (masl)	dg/dZ ($\mu\text{Gal}/\text{mm}$)	1996 (μGal)	2016 (μGal)	2018 (μGal)	Uncertainty (μGal)
CHA	Instituto de Biología UNAM, Chamela, Jal.	19°29'55.8"	105°02'41.8"	100	-3.709	978593393	978593377.4	978593371.3	1.9
MAN	CUNIVO, Universidad de Colima (Manzanillo)	19°07'27.4"	104°24'04.4"	18	-3.023	978594708	978594731.8	978594713.8	1.9
UGG	Instituto de Astronomía, U. de G., Guadalajara	20°40'29.7"	103°23'03.4"	1583	-2.80	978197026	978196995.6	978196979.8	3.8
UGP	Universidad. de Guadalajara, Campus Puerto Vallarta	20°42'21.1"	105°13'14.9"	13	-3.017	978589289	978589310.1	978589315.7	2.0
TEP	CICESE-UT3, Tepic, Nayarit	21°28'54.4"	104°50'58.0"	948	-3.00	N/M	978460630.9	978460630.4	1.9
AGS	INEGI-Aguascalientes	21°51'26.6"	102°17'04.05"	1888	-3.00	NA	978174962.0	N/M	1.9
IGU	Instituto de Geofísica, UNAM, Cd. de México	19°19'36.85"	99°10'34.02"	2280	-2.479	977927005	977926980.0	N/M	2.0
CNM	CENAM, Querétaro	20°32'17.9"	100°15'35.84"	1922	-3.00	N/M	978081416.0	978081416.8	1.2
CGE	CGEO-UNAM, Juriquilla,	20°42'7.2"	100°26'50.50"	1929	-3.00	N/M	978098243.0	978098242.9	1.8

THE JALISCO BLOCK (JB) RELEVANCE

The Jalisco Block (JB) is named after the western Mexico continental block (Figure 4a) about $40,000 \text{ km}^2$ of surface limited by three extensional zones in land, namely the Tepic-Zacoalco rift (TZG), the Colima graben (CG), and the Banderas graben (BG); seawards the JB is limited by the northernmost segment of the Meso-American trench. The JB is a continental micro-plate that moves independently respect to the Rivera plate (RP), North American plate (NAP), and the Michoacan Block (MB) (Luhr *et al.*, 1985; Allan, 1985; Johnson & Harrison, 1990; Bandy y

Pardo, 1994; Selvans *et al.*, 2010). The subduction rate of the RP under the JB is between 2.0 and 2.3 cm/yr (Nixon, 1982; Ferrari *et al.*, 2012), however, at the boundary of the Rivera and Cocos plates (CP) the rate of subduction increases to 5.8 cm/yr (Johnson and Harrison, 1990) or 3.8 cm/yr according to other estimations (Ferrari *et al.*, 2012). In any case, this difference in subduction speed must be accommodated through the Rivera transform fault (RFZ) and is highly probable that the Colima rift reflects in surface the divergence in speed and subduction angle of RP and CP at depth. Indeed, the RFZ could be extended in land to be correlated with the southern Colima graben (Alvarez and Yutsis, 2015A).




Figure 4. a) Location of AG stations (yellow circles) in the context of the Jalisco Block (JB) tectonic setting (compiled after Blatter and Hammersley, 2010, Rosas-Elguera *et al.*, 1996; Garduño *et al.*, 1998). Blue arrows are horizontal displacements (EARTHSCOPE-PBO, Plate Vel.), red squares are main cities. Latest Miocene to Quaternary faults shown with black lines. Closed red contours mark approximate extension of volcanic complexes surrounding the JB. CG, TZG and BG are Colima, Tepic-Zacoalco, and Banderas grabens. Gray arrows indicate convergence direction of tectonic plates (NAP=North America Plate, RP= Rivera Plate). b) and c) show the Jalisco Block batholith as imaged by the vertical gravity gradient (VGG) of the residual of Free Air anomaly (Basset and Watts, 2015). The dashed yellow contour highlights higher VGG's surrounding an apparently fragmented denser crustal block. Green arrows represent the magnitude and direction of cumulated vertical displacements respect to 1996 first AG measurements.

AG DATA COMPARISON

The differences in gravity for the periods 1996 to 2016, 1996 to 2018, and 2016 to 2018 are shown in Table 2 the yearly averages corresponding to these periods. The AG differences of the 2016 (left) and 2018 (right) results with respect to the 1996 measurements are shown schematically in Figure 4b. Green arrows at JB gravity stations (yellow circles) indicate the direction of displacement according to the sign of g . The figure shows that Puerto Vallarta (UGP)

and Manzanillo (MAN) stations have similar positive differences of g (+21 and +23 μGal s respectively). In contrast, at Chamela (CHA) and Guadalajara (UGG) stations a net negative difference of g is obtained, -15 and -29 μGal respectively. The background map in the Figure 4b is the vertical gravity gradient of the residual (VGG) of the FA anomaly from Basset and Watts (2015). The JB batholith is outlined with a dotted yellow line in the figure based upon the higher gravity anomalies. With only the 2016 set of g values (left) it is tempting to conclude that the reason why UGP and MAN stations are subsiding at about the same rate is because both are located within extensional zones (Bahía de Banderas and Colima rifts). However, the 2018 set of AG revealed that although the argument may hold for UGP station, which continued the subsiding rate, for CHA station it seems something more complicated as it changed from a subsiding regime to an uplifting regime in only two years. In contrast, stations CHA and UGG indicate uplift, which for the last two years increased two to three times above the yearly average (see Table 2).

Table 2.- Average differences in AG and yearly rate averages for the 20 years period (2016-1996), the 22 years period (2018-1996), and the later AG measurements (2018-2016).

STN	Z (masl)	Δg		Δg		Δg	
		2016-1996 μGal	Avg Δg $\mu\text{Gal/yr}$	2018-1996 μGal	Avg Δg $\mu\text{Gal/yr}$	2018-2016 μGal	Avg Δg $\mu\text{Gal/yr}$
CHA	100	-15.61	-0.78	-21.72	-0.99	-6.11	-3.06
MAN	18	23.83	1.19	5.77	0.26	-18.06	-9.03
UGG	1583	-30.37	-1.52	-46.19	-2.10	-15.82	-7.91
UGP	13	21.08	1.05	26.70	1.21	5.62	2.81
IGU	2280	-25.00	-1.25				
TEP	948					-0.50	-0.25
CNM	1929					0.76	0.38
CGE	1922					-0.09	-0.05

DISCUSSION OF RESULTS

Although the data base is still limited by the number of stations on the JB and the time series measurements of g , some general observations and conclusions can be drawn. The results obtained from the second campaign in 2018 proved that larger variations can occur within short periods of time. The observed differences between the 2016 and 2018 AG values were larger than expected. In the two years period the four gravity stations around the JB underwent important changes in the gravity field. The lowest change (+5.62 μGal) occurred at the station UGP, in Bahía de Banderas, where the positive sign implies sinking of the area in this short period. The highest change (-18.06 μGal) occurred at MAN station, within the Manzanillo Bay, equivalent to an uplift of 54.18 cm (Table 3) for the two years period assuming that the observed differences in gravity are all due to change in geoidal height. At Guadalajara city UGG station, the trend of g remained and confirmed uplift at this station with the second larger change observed (-15.82 μGal), which under the same assumption is equivalent to an uplift of 47.46 cm in two years. The remaining JB station (CHA) underwent the third larger uplift (-6.11 μGal) and remained consistent with a positive vertical displacement at a rate of 18.33 cm in the two years' period.




Figure 5. a) Map of western Mexico showing the configuration of ocean floor along the northernmost sector of the trench. The epicenters of the two large earthquakes of September 2017 are shown with red stars, green arrows indicate convergence direction, and shaded grey zones represent the extension of the trans Mexican volcanic belt (TMVB). Lighter grey area delineates younger volcanism. Yellow circles mark the location of the measured AG stations. b) Total magnetic anomaly map (Bankey *et al.*, 2002) of the JB and offshore magnetic records of the sea floor spreading (dotted lines). The Rivera and Tamayo transform fault zones (RFZ and TFZ respectively) are highlighted with a NW-SE continuous lines drawn along transverse low magnetic anomalies, coincident with the bathymetric traces of the transform fault under the ocean layer. The red arrow indicates the segment of the RP that experienced aseismic slip below the JB. c) magnetotelluric (MT) profile parallel to the coast at the mouth of southern Colima graben. JB is Jalisco block, MB Michoacan block, RP and CP are Rivera and Cocos plates. EGG is the Gordo graben, RFZ the Rivera transverse fault, which is apparently related to a central resistivity deep seated body.

In the middle of September 2017, two large earthquakes occurred in Mexico (Figure 5a) that were registered at some of UNAVCO's GPS stations (inset) at the location of the AG sites MAN, UGG and CHA in the JB. The recorded time series previous and after the two large earthquakes unveiled pre- and post-seismic vertical displacements even though the rupture occurred more than 500 km east of the JB. The earthquakes of magnitudes 8.2 and 7.1 with

epicenters in Chiapas (07/Sep/2017) and Puebla (19/Sep/2017) had lower crustal focal depths of 58 and 57 km respectively. Pre and post seismic activity after a large earthquake have been recorded within the JB previously by Hutton *et al.* (2001). The authors demonstrated using a local network of GPS stations that post seismic surface displacements can last for months, even years, and could be explained as consequence of aseismic slip subduction. The comparison of the AG data, measured before and after the two large September 2017 earthquakes, provided further evidence of important short-term variations in geoidal height in the JB (see Table 3).

Table 3. Average differences in AG and equivalent vertical displacement estimated using available gravity gradients at the gravity stations and assuming observed gravity differences are due solely to a change in height.

STN	Z (masl)	dg/dZ $\mu\text{Gal}/\text{cm}$	Avg Δg 2016-1996 $\mu\text{Gal}/\text{yr}$	Avg dZ cm/yr	Avg Δg 2018-1996 $\mu\text{Gal}/\text{yr}$	Avg dZ cm/yr	Avg Δg 2018-2016 $\mu\text{Gal}/\text{yr}$	Avg. dZ cm/yr
CHA	100	-3.709	-0.78	2.89	-0.99	3.67	-3.06	11.34
MAN	18	-3.023	1.19	-3.59	0.26	-0.78	-9.03	27.29
UGG	1583	-2.80	-1.52	4.25	-2.10	5.88	-7.91	22.14
UGP	13	-3.017	1.05	-3.16	1.21	-3.65	2.81	-8.47
TEP	948	-3.000					-0.50	-0.25
IGU	2280	-2.479	-1.25	3.09				

The unexpected rapid uplift at stations MAN (54.6 cm), CHA (22.7 cm), and UGG (44.3 cm), as well as the continued subsidence regime at station UGP (-17 cm), and the relative stability at station TEP (-0.5 cm) suggest us that the earthquake with epicenter in Puebla (M7.1) triggered the aseismic slip subduction of only a portion of the Rivera Plate. A seismic waveguide (Burg *et al.*, 1951) traveling eastwards from the earthquake source along the southern edge of the TMVB can account for the slow-slip activation of subduction in the JB region. Figure 5b shows the magnetic anomaly (Bankey *et al.*, 2002) of the JB and offshore magnetic records (dotted NS lines) of the sea floor spreading. The Rivera transform fault zone (RFZ) is highlighted with a NW-SE continuous line drawn along the transverse low magnetic anomaly that coincides with the bathymetric trace of the transform fault under the ocean layer.

According to these results, it appears that only a segment of the RP underwent aseismic slip under the JB (red arrow), otherwise gravity stations UGP and TEP should have also register important uplift as on stations CHA, UGG, and MAN. Aseismic slip is facilitated by a wet oceanic crust and likely a sizable amount of saturated subducted sediments. The relatively low resistivities (<250 Ohm-m) of the RP under the otherwise resistive continental crust (> 1,000 Ohm-m) is imaged along a magnetotelluric (MT) section in Figure 5c (Alvarez *et al.*, 2010). The electric structure along the profile (see location in a and b) suggest an irregular resistive crust of variable thickness at the NW sector of the profile (left of the image) associated to the JB micro plate. In the opposite side of the MT image it appears what may be the SW edge of the resistive Michoacan block (MB), separated from the JB by a relatively high conductivity region (~200 Ohm-m) with a central high resistivity (>2,000 Ohm-m) peak embedded. The extrapolation of the RFZ to the MT profile suggest that the central anomalous resistive body is related to the buried fault zone, which appears to be dipping northwestwards. The origin of the very low resistivity zone (< 20 Ohm-m) mediating between the JB and the RFZ is until now of unknown origin. In the context of the sea floor spreading and marine hydro-genetically precipitated ferromagnesian crust (Canet *et al.*, 2008) of the oceanic RP, the observed high conductivities can be associated to subducted conductive sea floor. However, concentration of mineralized fluids

can also account for the anomalous conductivity observed in the region (e.g. Corbo *et al.*, 2013), and its presence along the Rivera transform fault zone would imply low frictional contact surface.

CONCLUSIONS

Accurate and stable absolute gravity measurements are now possible in Mexico with the certified FG5X-252 absolute gravimeter of CENAM, which has become also a national metrological reference. International comparisons have been successful in this goal and are planned to continue. The instrument certification allowed us to set up of new AG stations in central and western Mexico and to repeat measurements at the Jalisco Block AG sites established by NOAA in 1996.

The repeated measurement campaigns of 2016 and 2018 revealed unusual variations of AG in the JB that we associate with an event of aseismic subduction of the RP that occurred within of the two years period. The large earthquake of the 19 September 2017 with epicenter in the State of Puebla is thought to have triggered a slow-slip event that uplifted stations CHA (22.3 cm), MAN (54.6 cm), and UGG (44.3 cm). TEP (0.25 cm) and UGP (-16 cm) stations were unaffected by the uplift registered at the other JB sites. These constraints allowed us to advance the hypothesis that only a segment of the RP moved under the JB, which was facilitated by a lateral displacement along the nearby RFZ transform fault. Packed fluids on top of the subducted oceanic sediments may play a key role in this process. Monitoring measurements at established benchmark sites in the JB showed that rapid changes in geodetic height can be triggered by large earthquakes occurring hundreds of km away. A waveguide along the southern edge of the TMVB is assumed to have started the low-frictional mechanism of aseismic subduction in the region.

ACKNOWLEDGMENTS

We acknowledge the financial support from PAPIIT, project IN116816, to carry out the JB measurements. We are grateful to Josefina Jacobo and Gina Villalobos (UT3-CICESE in Tepic), to Araceli Zamora of the University of Guadalajara, campus Puerto Vallarta, to Norma Barocio and Jorge Vega (Chamela station, IB-UNAM), to Ramón Sosa and colleagues (CUNIVO, UC-Manzanillo), to the authorities of the Institute of Astronomy of the U. of Guadalajara, and to Raúl Gómez from the INEGI in Aguascalientes for providing facilities during the AG acquisition campaigns. We greatly appreciate the assistance of Calixto Morales and Fernando Martínez during the measuring campaigns in the JB. Particularly, we would like to thank Dan Winester of NOAA, and Roger Billham from the University of Colorado at Boulder who kindly sheared their unpublished data. PTB Braunschweig funded logistical costs (travelling and freight shipping) of the LUH team and instrumentation to Mexico. We thank the anonymous referees that helped to improve this manuscript.

REFERENCES

- Allan, J. F. (1985). Sediment depth in the northern Colima graben from 3-D interpretation of gravity. *Geofísica Internacional*, 24(1).
- Alvarez, R. and Yutsis, V., 2015a. The elusive Rivera-Cocos plate boundary: not diffuse. From: Wright, T. J., Ayele, A., Ferguson, D. J., Kidane, T. & Vye-Brown, C. (eds) *Magmatic Rifting and Active Volcanism*. Geological Society, London, Special Publications, 420, 83-103. <http://doi.org/10.1144/SP420.8>
- Alvarez R., Arzate-Flores J.A., Corbo-Camargo F., 2010. The shape of the truncated, subducting oceanic slab at the terminus of the Middle America Trench. Annual Meeting of the American Geophysical Union, San Francisco, Cal.
- Bandy, W., & Pardo, M. (1994). Statistical examination of the existence and relative motion of the Jalisco and southern Mexico blocks. *Tectonics*, 13(4), 755-768.
- Bankey V., Cuevas A., Daniels D., Finn C.A., Hernandez I., Hill P., Kucks R., Miles W., Pilkington M., Roberts C., Roest W., Rystrom V., Shearer S., Snyder S., Sweeney R., Velez J., Phillips J.D., Ravat, D., 2002. Digital data grids for the magnetic anomaly map of North America. *U.S. Geological Survey, Open-File Report*, 02-414.
- Basset D. and Watts A.B., 2015. Gravity anomalies, crustal structure, and seismicity at the subduction zones: 2. Interrelationships between fore-arc structure and seismogenic behavior. *Geochemistry, Geophysics, Geosystems*, v16, 5
- Battler D.L. and Hammersley L., 2010. Impact of the Orozco fracture zone on the central Mexican Volcanic Belt. *J. of Volcanology and Geothermal Res.*, 197, 67-84
- Burg K.E., Ewing M., Press F., and Stulken J., 1951. A seismic wave guide phenomenon. *Geophysics*, 16(4), 594-612
- Canet Ch., Prol-Ledezma R.M., Bandy W.L., Schaaff P. Linares C., Camprubi A., Tauler E., and Mortera-Gutiérrez C., 2008. Mineralogical and geochemical constraints on the origin of ferromagnese crusts from the Rivera Plate (western margin Mexico). *Marine Geology*, 251, 47-59
- Corbo Camargo F., Arzate Flores J.A., Alvarez Béjar R., Aranda-Gómez J.J., y Vsevolod Yutsis, 2013. Subduction of the Rivera Plate beneath the Jalisco Block as imaged by magnetotelluric data. *Revista Mexicana de Ciencias Geológicas*, v. 30, n. 2, p. 268-281
- Ferrari, L., Orozco-Esquivel, T., Manea, V., & Manea, M., 2012. The dynamic history of the Trans-Mexican Volcanic Belt and the Mexico subduction zone. *Tectonophysics*, 522, 122-149.
- Francis, O., Baumann, H., Ullrich, C., Castelein, S., Van Camp, M., de Sousa, M.A., Melhorato, R.L., Li, C., Xu, J., Su, D., Wu, S., Hu, H., Wu, K., Li, G., Li, Z., Hsieh, W.C., Pálincás, V., Kostecký, J., Mäkinen, J., Näränen, J., Merlet, S., Pereira Dos Santos, F., Gillot, P., Hinderer, J., Bernard, J.D., Le Moigne, N., Fores, B., Gitlein, O., Schilling, M., Falk, R., Wilmes, H., Germak, A., Biolcati, E., Origlia, C., Iacovone, D., Baccaro, F., Mizushima, S., De Plaen, R., Klein, G., Seil, M., Radinovic, R., Sekowski, M., Dykowski, P., Choi, I.M., Kim, M.S., Borreguero, A., Sainz-Maza, S., Calvo, M., Engfeldt, A., Ágren, J., Reudink, R., Eckl, M., van Westrum, D., Billson, R., Ellis, B., 2015. CCM.G-K2 key comparison. *Metrologia* 52(1A):07009. doi:10.1088/0026-1394/52/1a/07009
- Garduño-Monroy, V. H., Saucedo-Girón, R., Jiménez, Z., Gavilanes-Ruiz, J. C., Cortes-Cortés, A., & Uribe-Cifuentes, R. M., 1998. La Falla Tamazula, límite suroriental del bloque Jalisco, y sus relaciones con el complejo volcánico de Colima, México. *Revista Mexicana de Ciencias Geológicas*, 15(2), 132-144.
- Hutton W., DeMets C., Sánchez O., Suárez G., and Stock J., 2001. Slip kinematics and dynamics during and after the 1995 October 9 Mw = 8.0 Colima-Jalisco earthquake, Mexico, from GPS geodetic constraints. *Geophys. J. Int.*, 146, 637-658
- IAG (1983). Resolutions adopted by the International Association of Geodesy. IUGG XVIII General Assembly, (https://iag.dgfi.tum.de/fileadmin/IAG-docs/IAG_Resolutions_1983.pdf, downloaded: 2015/10/18).
- Jiang, Z., Pálincás, V., Arias, F.E., Liard, J., Merlet, S., Wilmes, H., Vitushkin, L., Robertsson, L., Tisserand, L., Pereira Dos Santos, F., Bodart, Q., Falk, R., Baumann, H., Mizushima, S., Mäkinen, J., Bilker-Koivula, M., Lee, C., Choi, I.M., Karaboce, B., Ji, W., Wu, Q., Ruess, D., Ullrich, C., Kostecký, J., Schmerge, D., Eckl, M., Timmen, L., Le Moigne, N., Bayer, R., Olszak, T., Ágren, J., Del Negro, C., Greco, F., Diamant, M., Deroussi, S., Bonvalot, S., Krynski, J., Sekowski, M., Hu, H., Wang, L.J., Svitlov, S., Germak, A., Francis, O., Becker, M., Inglis, D., Robinson,

- I., 2012. The 8th International Comparison of Absolute Gravimeters 2009: the first Key Comparison (CCM.G-K1) in the field of absolute gravimetry. *Metrologia* 49, 666–684, doi:10.1088/0026-1394/49/6/666.
- Johnson, C. A., & Harrison, C. G. A., 1990. Neotectonics in central Mexico. *Physics of the Earth and Planetary Interiors*, 64(2-4), 187-210.
- Luhr, J. F., Nelson, S. A., Allan, J. F., & Carmichael, I. S., 1985. Active rifting in southwestern Mexico: Manifestations of an incipient eastward spreading-ridge jump. *Geology*, 13(1), 54-57.
- Niebauer, T. M., Sasagawa, G. S., Faller, J. E., Hilt, R., and Klopping F., 1995. A new generation of absolute gravimeters, *Metrologia*, 32, 159-180.
- Niebauer, T. M., Billson, R., Schiel, A., van Westrum, D., and Klopping F., 2013. The self-attraction correction for the FG5X absolute gravity meter, *Metrologia*, 50, 1-8.
- Nixon, G. T., 1982. The relationship between Quaternary volcanism in central Mexico and the seismicity and structure of subducted ocean lithosphere. *Geological Society of America Bulletin*, 93(6), 514-523.
- Ramírez-Herrera, T., Kostoglodov, V., and Urrutia-Fucugauchi, J., 2004. Holocene-emerged notches and tectonic uplift along the Jalisco coast, Southwest Mexico. *Geomorphology* 58 291–58 304. Elsevier
- Ramírez-Herrera, T., Kostoglodov, V., and Urrutia-Fucugauchi, J., 2010. Overview of recent Coastal Tectonic Deformation in the Mexican Subduction Zone. *Pure and Applied Geophysics*. Springer International Publishing
- Rosas-Elguera, J., Ferrari, L., Garduño-Monroy, V. H., & Urrutia-Fucugauchi, J., 1996. Continental boundaries of the Jalisco block and their influence in the Pliocene-Quaternary kinematics of western Mexico. *Geology*, 24(10), 921-924.
- Rothleitner Ch., Niebauer T.M., and Francis O., 2014. Measurement of the speed-of-light perturbation of free-fall absolute gravimeters. *Metrologia*, 51, Number 3 <https://iopscience.iop.org/article/10.1088/0026-1394/51/3/L9/pdf>
- Selvans, M. M., Stock J.M., DeMets Ch., Sanchez O., and Márquez-Azúa B., 2010. Constraints on Jalisco Block Motion and Tectonics of the Guadalajara Triple Junction from 1998-2001 Campaign GPS Data. *Pure and Applied Geophysics*, 168(8–9), pp. 1435–1447. doi: 10.1007/s00024-010-0201-2.
- Schilling, M., and Timmen, L., 2016. Traceability of the Hannover FG5X-220 to the SI Units. International Association of Geodesy Symposia, DOI 10.1007/1345_2016_226, *Springer International Publishing*
- Tamura, Y., 1987. A harmonic development of the tide generating potential. *Bull. d'Inf., Marées Terrestres*, 99, 6813–6855, Bruxelles
- Timmen, L., Engfeldt, A., and Scherneck, H.G., 2015. Observed secular gravity trend at Onsala station with the FG5 gravimeter from Hannover. *J. of Geod. Sc.*, 5:18-25, DeGruyter Open, DOI 10.1515/jogs-2015-0001
- Timmen, L., and Wenzel, H.-G., 1995. Worldwide synthetic gravity tide parameters. In: Sünkel, H., and Marson, I. (eds), *Gravity and Geoid*. Proceedings of IAG Symposium, 113, *Springer*, Berlin, Heidelberg, pp. 92–101
- Wahr J., 1985. Deformation induced by polar motion. *J. of Geophys. Res.* 92(B2), p. 1281-128 *Geochemistry* 21 (2006) 1064–1072.