Phytoplankton biomasses and hydrographic conditions during El Niño 1997-1998 in Bahía Concepción, Gulf of California, Mexico

David J. López-Cortés¹, José J. Bustillos-Guzmán¹, Ismael Gárate-Lizárraga², Francisco E. Hernández-Sandoval¹ and Iban Murillo-Murillo¹

¹Centro de Investigaciones Biológicas del Noroeste (CIBNOR), La Paz, Baja California Sur, México

Received: September 3, 2000; accepted: May 20, 2002

RESUMEN

Los cambios en las condiciones físico-químicas y los pigmentos fitoplanctónicos, en Bahía Concepción, Golfo de California, se analizan para el periodo 1997 a 1999 con el objetivo de observar el efecto de El Niño 1997-1998 en esta bahía. Tres diferencias notables en los parámetros medidos ocurrieron durante 1998: Temperaturas mayores en la columna de agua, un retrazo de 2-3 meses en el inicio del período de la estratificación y concentraciones menores de nutrientes (nitratos y fosfatos) y peridinina. Estos datos sugieren la presencia de Agua Ecuatorial Superficial (altas temperaturas y baja concentración de nutrientes) con un efecto negativo en la proliferación de dinoflagelados. Se discuten las posibles causas de la baja en la concentración de los dinoflagelados.

PALABRAS CLAVE: El Niño, huellas pigmentarias, Bahía Concepción, Golfo de California.

ABSTRACT

The biomass of phytoplankton groups, particularly dinoflagellates as indicated by their signature pigment peridinin, and the physicochemical conditions in Bahía Concepción, Gulf of California are analyzed for the years 1997-1999, in order to observe the changes in phytoplankton and hydrographic structure linked to El Niño 1997-98. Three significant changes occurred in 1998 compared to 1997 and 1999: higher temperatures in the water column, lower nutrient and peridinin concentrations, and a 2 to 3 month delay of the stratification period. These data suggest the influence of Equatorial Surface Water (low nutrients, high temperatures) and a negative effect on the proliferation of dinoflagellate groups during El Niño 1997-98. Possible causes of the low dinoflagellate biomass are discussed.

KEY WORDS: El Niño, pigment signatures, Bahía Concepción, Gulf of California.

INTRODUCTION

In general, the El Niño effects along the American west coast consist of a relaxation of upwelling, a decrease of nutrients, and a decline of primary and secondary productivities (Lynn et al., 1998). A decrease in phytoplankton biomass along the coasts of Chile and Peru (Cowles et al., 1977; Barber et al., 1983; Guillén et al., 1985; Avaria and Muñoz, 1987) and the western coast of Baja California (Torres-Moye and Álvarez-Borrego, 1985; Martínez-López, 1993; Gárate-Lizárraga and Siqueiros-Beltrones, 1998) have been described. In the Gulf of California, the consequences appear to be different. The primary productivity and phytoplankton and zooplankton biomasses increased during El Niño 1982-83 (Baumgartner et al., 1985; Valdéz-Holguín and Lara-Lara, 1987; Bustillos-Guzmán, 1990; Lavaniegos-Espejo and Lara-Lara, 1990; Álvarez Borrego and Lara-Lara, 1991). This response was attributed to a lower grazing pressure and to a higher incidence of tropical and subtropical species during the El Niño (Valdéz-Holguín and Lara-Lara, 1987; Álvarez Borrego and Lara-Lara, 1991). A stronger advection of nutrient-rich subsurface subtropical waters into the Gulf and their transport to the surface by upwelling has also been used to explain the high productivity (Baumgartner and Christensen, 1985; Robles and Marinone, 1987). Thus, the Gulf of California acts as a chemostatic system, where the nutrient supply is provided by the horizontal subsurface flow of nutrients (Baumgartner et al., 1985). Studies based on monthly composites of coastal-zone color-scanner (CZCS) satellite imagery show that, in general, variability of phytoplankton biomass (as chlorophyll concentration) in the northern and central part of the Gulf of California is unrelated to El Niño (Santamaríadel Angel et al., 1994).

Phytoplankton in coastal lagoons around the Gulf of California is clearly influenced by Gulf waters (Gilmartin and Revelante, 1978; Martínez-López and Gárate-Lizárraga, 1997). Thus, an effect may also be expected during El Niño events, but long time-series data are unavailable as yet to test this hypothesis. For Bahía Mazatlán, near the mouth of the Gulf of California, a decrease in the frequency and du-

²Depto. de Plancton y Ecología Marina, Centro Interdisciplinario de Ciencias Marinas (CICIMAR-IPN), La Paz, Baja California Sur, México

ration of red tides has been noted (Cortés-Altamirano, 1987; Cortés-Altamirano *et al.*, 1995; Cortés-Altamirano and Alonso-Rodríguez, 1997). In this study, we analyze the physicochemical conditions and phytoplankton biomass of Bahía Concepción during El Niño 1997-98.

STUDY AREA

Bahía Concepción is on the west coast of the Gulf of California ((26° 33' - 26° 53' N; 111° 42' - 112° 56' W). It is approximately 45 km long by 10 km at the widest part. The depth in the central channel is 30 m (Ramírez-Guillén, 1983). From May to October the prevailing winds are weak and primarily from the south. In late fall through early spring, the winds are strong and from the north (Baqueiro-Cárdenas et al., 1978; Thunell et al., 1994). This bay has anti-estuarine features, and the mean annual surface temperature is 24.9 °C, with a lowest mean of 17.5°C (January) and a maximum mean of 32.1 °C (September). Salinity has an annual mean of 35.3 psu, with a minimum and maximum of 34.6 and 37 psu (Félix-Pico and Sánchez, 1976). The concentrations of dissolved oxygen range from 5.4 mL/L in Spring to 5.95 mL/L in Winter. However, at the bottom anoxic or hypoxic conditions have been observed (Gilmartin and Revelante, 1978; Lechuga-Devéze and Morquecho-Escamilla, 1998; Lechuga-Devéze et al., 2000).




Fig. 1. Location of the study area and sampling station in Bahía Concepción, Gulf of California.

MATERIAL AND METHODS

During 1997, 1998, and 1999 a sampling program was done at a fixed sampling station in the central part of Bahía Concepción (Figure 1). Water samples were obtained at 0, 5, 10, 15, 20, 25, and 27 m depth with a Van-Dorn bottle to determine temperature, dissolved oxygen, nitrates and phosphates, and photosynthetic pigments. Sampling intervals were 3 to 5 days in 1997 (February to May), 3 days in 1998 (February to November), and biweekly in 1999 (January to December).

Seawater temperature was measured with a bucket thermometer. Dissolved oxygen, nitrates, and phosphates were determined following methods described by Strickland and Parson (1972). Photosynthetic pigments were measured by HPLC (Vidussi *et al.*, 1996). Identification and quantification of pigments were made as described by Bustillos-Guzmán *et al.* (1995). The thermal stratification index (ST: °C/m) was computed as the difference of temperature m⁻¹ (0 to 27 m) calculated from each 5 m data. Homogenous water-column (ST=<0.05); Transition period (> 0.05 < 0.2); Stratified water column (> 0.2) (Bustillos-Guzmán *et al.*, 1995).

RESULTS

Temperature

A homogeneous water column with temperature of 18 °C was recorded in February 1997. At the end of March, temperature increased to 20 °C (Transition period). In May, the thermocline was completely developed between 10 and 20 m and the surface temperature reached 27 °C. A stratified water column is observed (Figure 2A). From January to May 1998, water column temperatures were vertically homogeneous (19 to 25 °C). From middle April to the end of June the first transition period was observed. In June, stratification was present again and it remained until September. During this period, water column temperatures were between 24 and 31 °C and the thermocline was found between 10- and 20-m depth. Surface temperature reached 31 °C (Figure 2B). Second transition period during 1998 was observed from middle September to early October.

From January to April 1999, the water column was homogeneous with temperatures between 16.0 and 21.5 °C. First transition period was observed from late April to late May. Water column stratification was detected during the first days of May. From June to September, the water column remained well-stratified with temperatures between 22.0 and 31.0 °C. Second transition period was observed from middle to end of September. Homogeneous water




Fig. 2. Variation of temperature (°C) during 1997 (A), 1998 (B), and 1999 (C) in Bahía Concepción, Gulf of California.




Fig. 3. Stratification index (ST) during 1997, 1998, and 1999 in Bahía Concepción, Gulf of California.

column conditions were again reached between October and December (19.0 and 28.0 °C) (Figure 2C).

The stratification index showed that during 1997 stratification was reached earlier. The delay period of about 2 months is evident in 1998. In 1999 stratification began in between the time of stratification in 1997 and 1998. The end of the stratification was similar in 1998 and 1999, therefore 1998 has a shorter stratification period (Figure 3).

Phosphates and nitrates

In general, values of phosphate and nitrate concentration increased with stratification, although lower values were clearly recorded in 1998 (Figure 4). The phosphate concentration was higher in 1999 when the value reached 30 mM/m², in contrast with the low values found in 1998 (< 10 mM/m² for the same period) (Figure 4A). Peak values on nitrates were around 20 mM/m² during 1998, whereas peak values for 1997 were 30 and for 1999 were 40 mM/m² (Figure 4B).

Phytoplankton pigments

In general, an increase of pigments occurred during the hydrographic transitional periods (Figure 5). Chlorophyll *a* values were similar form year to year, and with the exception of peak values (> 70 mg m⁻²) that occurred (3 for 1997, 1 for 1998, and 1 for 1999), they were lower than 40 mg m⁻² (Figure 5A). For peridinin concentrations, the outstanding feature is the low value during 1998 (in general lower than 1.5 mg/m²) and peak values found during July

1999 (Figure 5B) caused by a dinoflagellate bloom. Fucoxanthin was the most important carotenoid and closely followed the variation of the chlorophyll *a*, suggesting that diatoms are responsible for most of the phytoplankton biomass changes in the bay (Figure 5C). The chlorophyll *b*-containing groups (Chlorophyceae and Prasinophyceae) show no important differences from one year to the next (Figure 5D).

DISCUSSION

Hydrographic conditions were similar to those reported previously (Reyes-Salinas, 1994; Lechuga-Devéze and Morquecho-Escamilla, 1999). Dominance of diatoms and dinoflagellates is a community phytoplankton feature for Bahía Concepción (Martínez-López and Gárate-Lizárraga, 1994; 1997), as confirmed by the high concentration of fucoxanthin and peridinin (Figure 5). Additionally, the pigment signatures showed an important contribution of the chlorophyll b-containing groups. Warmer temperatures at depth as well as a time-delay in the stratification were noted during 1998. These differences could be caused by the normal annual and interannual surface temperature variability in the Gulf of California (Soto-Mardones et al., 1999) and the wind effect on the hydrographic patterns (Santa-María del Angel et al., 1994). These differences were also accompanied by a lower concentration of phosphate, nitrate, and chlorophyll a, suggesting that they can be linked to the El Niño 1997/1998 phenomenon. Let us explore this possibility.

Evidence of El Niño off the Mexican coastal zone, in the form of surface Tropical Pacific water (Durazo and




Fig. 4. Integrated values of phosphate (A) and nitrate (B) concentration during 1997, 1998, and 1999 in Bahía Concepción, Gulf of California.

Baumgartner, in press) was detected in June 1997 (Filonov and Tereshchenko, 2000). Positive anomalies of the sea level and sea surface temperature (SST) also linked to the El Niño (Baumgartner and Christensen, 1985) were recorded during July and August 1997 in the Gulf of California (Thunell *et al.*, 1999). SSTs in Bahía Concepción were similar for the three years (Figure 6A) but temperatures below 15 meters were higher during 1998. This is better shown when comparing the mean water column temperature (Figure 6B). Therefore, it is plausible to assume that the warming effect of El Niño was present in Bahía Concepción until October 1998 and began during the second half of 1997. Positive SST anomalies were also recorded in the California waters (Lynn *et al.*, 1998).

Transitional hydrographic periods are important for phytoplankton because an enhancement of their biomass occurs because the increase of the nutrients (Kiørboe, 1993; Townsend *et al.*, 1994; Bustillos-Guzmán *et al.*, 1995; Thunell *et al.*, 1996). These features were recorded in Bahía

Concepción, but in 1998 the beginning of the stratification period was delayed for about 3 months (Figure 3). According to Álvarez-Borrego and Lara-Lara (1991), stratification in the Gulf of California begins when the northwest winds diminish. It is plausible that a prolongation of northwest winds was the cause of such a delay (unpublished data).

The stratification period in Bahía Concepción is also important because there is a high organic-matter oxidation and nutrient production in the bottom waters. Lechuga-Devéze *et al.* (2000) calculated that for a "normal" 225 day stratification period, N-Nitrate released by mineralization of organic matter is 68 mmol N-Nitrate m⁻²y⁻¹. Thus, using a stratification period of only 135 days (for 1998), only 40.8 mmol N-Nitrate m⁻²y⁻¹ should be produced (40% less). The shorter mineralization time during 1998 may be linked to the lower NO₃ and PO₄ concentrations.

During the transition period from a well-mixed to a stratified water column, nutrient-rich waters, coming in from




Fig. 5. Integrated values of pigment concentration during 1997, 1998, and 1999 in Bahía Concepción, Gulf of California. A) Chlorophyll *a.* B) Peridinin. C) Fucoxanthin. D) Chlorophyll *b.*




Fig. 6. Sea surface temperature (SST) (A) and mean temperature (B) during 1997, 1998, and 1999 in Bahía Concepción, Gulf of California.

the Gulf of California, and nitrate and nitrite production inside the bay provide an optimum environment for phytoplankton, mainly diatoms and dinoflagellates (Martínez-López and Gárate-Lizárraga, 1997). The association between this transitional period with high nutrient and pigment concentrations is clearly observed in our data. According to Álvarez-Borrego and Schwartzlose (1979), during spring and summer the central part of the Gulf of California is influenced on the surface mainly by the Equatorial Surface Water (ESW), characterized by low salinity (<35.0 psu) and low nutrient concentrations. Additionally, the ESW flow into the Gulf is intensified and remains for longer periods when the El Niño is present (Torres-Orozco, 1993; Castro et al., 2000; Lavin, 2002). Flow intensification and a longer permanence of ESW in the area may also be the cause of the low nutrients recorded during 1998.

Important blooms of dinoflagellates have been recorded in Mazatlán, Guaymas, and Bahía de La Paz during the upwelling periods (Cortés-Altamirano *et al.*, 1995; Gárate-Lizárraga *et al.*, 2001). But the decrease in the intensity of upwelling that occurred during El Niño

conditions (Santamaría-del Angel *et al.*, 1994) has a negative effect on the dinoflagellate blooms (Cortés-Altamirano 1987; Manrique and Molina, 1997). According to the above results and our data (high temperatures and the low nutrient concentration during 1998), it is plausible to assume that the 1997-1998 El Niño caused environmental conditions that negatively affected the dinoflagellate group. Phytoplankton biomass was reestablished in 1999 together with hydrographic conditions.

The main dinoflagellate blooming species in Bahía Concepción is *Noctiluca scintillans*, and toxic species from the genera *Alexandrium*, *Dinophysis*, *Prorocentrum*, and *Gymnodinium* have been also reported (Gárate-Lizárraga, 1991; Lechuga-Devéze and Morquecho-Escamilla, 1998; Gárate-Lizárraga *et al.*, 2001). The highest dinoflagellate concentrations occurred during the mixed-stratified hydrographic period under a narrow temperature window of 23 to 26 °C (Lechuga-Devéze and Morquecho-Escamilla, 1998; Lechuga-Devéze *et al.*, 2000). In 1998, temperatures were in the upper limit of this range and therefore could be linked to the low dinoflagellate biomass. Two dinoflagellate species bloomed during 1999 in Bahía Concepción,

Gymnodinium catenatum (May) and Alexandrium affine (July) (Gárate-Lizárraga et al., 2002). These species were responsible for the elevated peridinin concentration (Figure 6B). Although the SST was high, both species were located below the thermocline, mainly above the oxycline, where temperature is lower (22 to 24 °C). Nutrients could be supplied by a diffusion mechanism from the anoxic zone as suggested by Bustillos-Guzmán et al. (2000).

CONCLUDING REMARKS

Differences in magnitude as well as environmental conditions previous and post the El Niño event make difficult a direct comparison between an event and other. This hold particularly true when time series data are not available as is our case. However, our data set suggest that conditions present in Bahía Concepción during the El Niño 1997-1998, did not favored the proliferation of dinoflagellates because the high temperatures and low nutrient conditions. This effect, however, have to be corroborated with longer timeseries data.

ACKNOWLEDGEMENTS

This project was financed by Conacyt (007P Ñ-1297; 33684-V) and CIBNOR (project AYCG11, GEA-3). Additional support came from the IPN (Project: DEGEPI 990318. I.G.L. is a COFAA, EDI, and Conacyt fellow (Beca 138138). Suggestions and comments by R. Durazo and an external reviewer greatly improved the paper. We thank Dr. Ellis Glazier for editing the English-language text, and an anonymous referee.

BIBLIOGRAPHY

- ÁLVAREZ-BORREGO, S. and R. A. SCHWARTZLOSE, 1979. Masas de agua del Golfo de California. *Cien. Mar.*, 6(2), 43-63
- ÁLVAREZ-BORREGO, S. and J. R. LARA-LARA, 1991. The physical environment and primary Productivity of the Gulf of California. *In*: Simoneit, B.R.T. and Drophin, JP. (eds.). The Gulf and Peninsular Province of the Californias. Am. Assoc. Petr. Geol. Memoir. 47, 555-567.
- AVARIA, S. and P. MUÑOZ, 1987. Effects of the 1982-1983 El Niño on the marine phytoplankton off northern Chile. *J. Geophys. Res.*, 92, 14,369-14,382.
- BAQUEIRO, C. E., J. A. MASSO and A. VÉLEZ. 1983. Crecimiento y reproducción de una población de caracol chino *Hexaplex erythristomus*, (Swainson, 1831) de Bahía Concepción, B.C.S. *Cien. Pesq.*, 4, 19-31.

- BARBER, R. T., S. ZUTA, J. KOGELSCHATZ and F. CHÁVEZ, 1983. Temperature and nutrient condition in the equatorial Pacific, October 1982. *Trop. Oc. Atmos. News* 16, 15-17.
- BAUMGARTNER, T. and N. CHRISTENSEN, 1985. Coupling of the Gulf of California to large-scale climatic variability. *J. Mar. Res.*, *43*, 825-848.
- BAUMGARTNER, T., V. FERREIRA-BARTRINA, H. SHRADER and A. SOUTAR, 1985. A 20 year varve record of siliceous phytoplankton variability in the central Gulf of California. *Mar. Geol.*, 64, 113-129.
- BUSTILLOS-GUZMÁN, J. J., 1990. Biomasa proteica, de carbohidratos y clorofila de las fracciones de nanopartículas y micropartículas de la región de las grandes islas y central del Golfo de California. Tesis de Maestría. CICIMAR-IPN., La Paz, México. 96 pp.
- BUSTILLOS-GUZMÁN, J., H. CLAUSTRE and J. C. MARTY, 1995. Specific phytoplankton signatures and their relationship to hydrographic conditions in the coastal northwestern Mediterranean Sea. *Mar. Ecol. Prog. Ser.*, 124, 247-258.
- BUSTILLOS-GUZMÁN, J., D.J. LÓPEZ-CORTÉS, F. HERNÁNDEZ and I. MURILLO, 2000. Pigment signatures associated with an anoxic coastal zone: Bahía Concepción, Gulf of California. *J. Exp. Mar. Biol. Ecol.*, 249, 77-88.
- CASTRO, R., A.S. MASCARENAS, R. DURAZO and C. A. COLLINS, 2000. Seasonal variation of the temperature and salinity at the entrance to the Gulf of California, Mexico. *Cien. Mar.*, 26(4), 561-583.
- CORTÉS-ALTAMIRANO, R., 1987. Observaciones de Mareas Rojas en la Bahía de Mazatlán, Sinaloa, México. *Cien. Mar.*, 13(4), 1-19.
- CORTÉS-ALTAMIRANO, R., D.U. HERNÁNDEZ-BECERRIL and R. LUNA-SORIA, 1995. Mareas rojas en México: una revisión. *Rev. Lat-Amer. Microbiol.*, 37, 343-352.
- CORTÉS-ALTAMIRANO, R. and R. ALONSO-RODRÍGUEZ, 1997. Mareas rojas durante 1997 en la bahía de Mazatlán, Sinaloa, México. *Cienc. Mar, UAS.* 15, 31-37.
- COWLES, T. J., R. T. BARBER and O. GUILLÉN, 1977. Biological consequences of the 1975 El Niño. *Science*, 195, 285-287.

- DURAZO, R. and T. BAUMGARTNER, (in press). Evolution of Oceanographyc conditions off Baja California: 1997-1999. *Prog. Oceanogr.*
- FÉLIX, P. E. and R. S. SÁNCHEZ, 1976. Tercer informe final del programa de orientación técnica para el aprovechamiento de los recursos naturales, existentes y prácticas de maricultivos en Bahía Concepción y Ensenada de La Paz. Secretaría de Recursos Hidraúlicos, 20 p.
- FILONOV, A. and L. TERESHCHENKO, 2000. El Niño 1997-1998 monitoring in mixed layer off the west coast of Mexico. *Geophys. Res. Lett.*, 27 (5), 705-711.
- GÁRATE-LIZÁRRAGA, I., 1991. Análisis de una marea roja causada por *Noctiluca scintillians* (Macrtney) Ehr. en Bahía Concepción Baja California Sur en febrero de 1989. *Rev. Invest. Cient.* 2(1):35-43.
- GÁRATE LIZÁRRAGA, I. and D. A. SIQUEIROS BELTRONES, 1998. Time variations in phytoplankton assemblages in a subtropical lagoon system after the 1982/83 El Niño event (1984/86). *Pac. Sci. 52 (1)*, 79-97.
- GÁRATE-LIZÁRRAGA, I., M. L. HERNÁNDEZ-OROZCO, C. BAND-SCHMIDT and G. SERRANO-CASILLAS, 2001. Red tides along of the coasts of the Baja California Peninsula, Mexico (1984 to 2001). *Oceán.*, 16(2), 127-134.
- GÁRATE-LIZÁRRAGA, I., D. A. LÓPEZ-CORTÉS, J. J. BUSTILLOS-GUZMÁN, F. E. HERNÁNDEZ-SANDOVAL and I. MURILLO-MURILLO, 2002. Physicochemical characteristics and phytoplankton biomass during El Niño 1997-1998 in Bahía Concepción, Gulf of California. International Symposium on North Pacific Transitional Areas. La Paz, Mexico. p. 42.
- GILMARTIN, M. and N. REVELANTE, 1978. The Phytoplankton characteristics of the Barrier Island Lagoons of the Gulf of California. *Est. Coast. Mar. Sci.*, 7, 29-47.
- GUILLÉN, O., N. LOSTAUNAU and M. JACINTO, 1985. Caractéristicas del fenómeno El Niño 1982-83. Pages 11-22 *In*: W. Arntz, A. Landa, and J. Tarazona (eds). El Niño, su impacto en la zona marina. Boletin del Instituto del Mar, Perú. Vol. Extraordinario.
- KIØRBOE, T., 1993. Turbulence, Phytoplankton Cell size, and the structure of Pelagic food webs. *Adv. Mar. Biol.*, 29, 1-72.

- LAVANIEGOS-ESPEJO, B. and R. LARA-LARA, 1990. Zooplankton of the Gulf of California after the 1982-1983 El Niño: Biomass distribution and abundance. *Pac. Sci.*, 18, 297-310.
- LAVIN, M. F., E. PALACIOS-HERNÁNDEZ and C. CA-BRERA, 2002. Sea surface temperature anomalies in the Gulf of California. *Geofis. Int.* (This issue).
- LECHUGA-DEVÉZE, C. and M. L. MORQUECHO-ESCAMILLA, 1999. Early spring harmful Phytoplankton in Bahía Concepción, Gulf of California. *Bull. Mar. Sci.*, 64, 1-10.
- LECHUGA-DEVÉZE, C., M. L. MORQUECHO-ESCAMI-LLA, A. REYES-SALINAS and J. R. HERNÁNDEZ-ALFONSO, 2000. Environmental natural disturbance at Bahía Concepción. Gulf of California. *In*: Munawar, M., Lawrence, S. G, Munawar, I. F. and Malley, D. F (Eds.) Ecosystem of Mexico: Status and scope. Backhuys Publishers B.V., Netherlands. pp. 245-255.
- LYNN, J. R., T. BAUMGARTNER, J. GARCÍA, C. A. COLLINS, T. L. HAYWARD, K. D. HYRENBACH, A. W. MANTYLA, T. MURPHREE, A. SHANKLE, F. B. SCHWING, K. M. SAKUMA and M. J. TEGNER, 1998. The State of the California Current, 1997-1998: Transition to El Niño conditions. *Cal.COFI. Rep.*, 39, 25-51.
- MANRIQUE, F. A. and R. E. MOLINA, 1997. Presencia de mareas rojas en la Bahía de Bacochibampo, Guaymas, Sonora, México. *Hidrobiol.*, 7, 81-84.
- MARTÍNEZ-LÓPEZ, A., 1993. Efectos del evento El Niño 1982-83 en la estructura del fitoplancton en la costa occidental de Baja California Sur. Tesis de Maestría, CICIMAR-IPN, La Paz, México. 83 p.
- MARTÍNEZ-LÓPEZ, A. and I. GÁRATE-LIZÁRRAGA, 1994. Cantidad y calidad de la materia orgánica particulada en Bahía Concepción en la temporada de reproducción de la almeja catarina *Argopecten circularis* (Sowerby, 1835). *Cien. Mar.*, 20 (3), 301-320.
- MARTÍNEZ-LÓPEZ, A. and I. GARATE-LIZÁRRAGA, 1997. Variación diurna de la materia orgánica particulada en una laguna costera del Golfo de California. *Rev. Biol. Trop.*, 45 (3), 1310-1317.
- RAMÍREZ-GUILLÉN, P. A., 1983. Sistemática, Ecología y Biogeografía de los crustáceos anomuros de Bahía Concepción, B. C. S., Tesis Profesional, U.A.N.L. 78 pp.

- REYES-SALINAS, A., 1994. Relación entre estructura hidrográfica y la abundancia, distribución y origen de diferentes expresiones de biomasa del seston orgánico en Bahía Concepción, Golfo de California. Tesis de Licenciatura, UNAM, Campus Iztacala, 53 p.
- ROBLES, J. M. and S. G. MARINONE, 1987. Seasonal and interannual thermo-haline variability in the Guaymas basin of the Gulf of California. *Cont. Shelf Res.*, 7, 715-733.
- SANTAMARÍA-DEL-ANGEL, E., S. ÁLVAREZ-BORRE-GO and F. E. MÜLLER-KARGER, 1994. The 1982-1984 El Niño in the Gulf of California as seen in coastal zone color scanner imagery. *J. Geophys. Res.*, *99(C4)*, 7423-7431.
- SOTO-MARDONES, L. S. G. MARINONE and A. PARÉS-SIERRA, 1999. Time and spatial variability of sea surface temperature in the Gulf of California. *Cien. Mar.*, 25 (1), 1-30.
- STRICKLAND, J. D. and R. PARSONS, 1972. A practical handbook of seawater analysis. *Fish. Res. Bd. Can. Bull.*, *167*, 207-211.
- TORRES-OROZCO, E, 1993. Análisis volumétrico de las masas de agua del Golfo de California, M.S. thesis, Cent. De Invest. Cient. y Educ. Super. de Ensenada, Baja California, México. 80 p.
- THUNELL, R. C., C. J. PRIDE, E. TAPPA and F. E MULLER-KARGER, 1994. Biogenic silica fluxes and accumulation rates in the Gulf of California. *Geology*, 22, 303-306.
- THUNELL, R., C. PRIDE, P. ZIVERI, F. MULLER-KARGER, C. SANCETTA and D. MURRAY, 1996. Plankton response to physical forcing in the Gulf of California. J. Plank. Res., 18 (11), 2017-2026.

- THUNELL, R., E. TAPPA, C. PRIDE and E. KINCAID, 1999. Sea-surface temperature anomalies associated with the 1997-1998 El Niño recorded in the oxygen isotope composition of planktonic Foraminifera. *Geology* 843-846.
- TORRES-MOYE, G. and S. ÁLVAREZ-BORREGO, 1985. Efectos de El Niño en los nutrientes y el fitoplancton de verano de 1983, en aguas costeras de Baja California occidental. *Cien. Mar.*, 11 (3), 107-113.
- TOWNSEND, D. W., L. M. CAMMEN, P.M. HOOLIGAN, D. E. CAMPBELL and N. R. PETTIGREW, 1994. Causes and consequences of variability in the timing of spring phytoplankton blooms. *Deep-Sea Res.*, *141*, 747-765.
- VALDÉZ-HOLGUÍN, J. E. and R. LARA-LARA, 1987. Productividad primaria en el Golfo de California: efectos del evento El Niño 1982-1983. *Cien. Mar.*, *3*, 34-50.
- VIDUSSI, F. H. CLAUSTRE, J. BUSTILLOS-GUZMÁN, C. CAILLEAU and J. C. MARTY, 1996. Rapid HPLC method for determination of phytoplankton chemotaxonomic pigments: separation of chlorophyll *a* from divinyl-chlorophyll *a* and zeaxanthin from lutein. *J. Plank. Res.* 18, 2377-2382.

David J. López-Cortés¹, José J. Bustillos-Guzmán¹, Ismael Gárate-Lizárraga², Francisco E. Hernández-Sandoval¹ and Iban Murillo-Murillo¹

¹Centro de Investigaciones Biológicas del Noroeste (CIBNOR), Apdo. Postal 128, La Paz, Baja California Sur, C.P. 23000, México

Email: dlopez@cibnor.mx

²Departamento de Plancton y Ecología Marina. Centro Interdisciplinario de Ciencias Marinas (CICIMAR-IPN). Apdo. Postal 592, La Paz, Baja California Sur, C.P. 23000, México